

## Supplementary File

<b>Supplementary Table 1: Prevalence of PEW among males and females according to various nutritional assessment tools</b>						
<b>Tool</b>	<b>Overall</b>		<b>Males (n=104)</b>		<b>Females (n=42)</b>	
	<b>Frequency</b>	<b>Percentage</b>	<b>Frequency</b>	<b>Percentage</b>	<b>Frequency</b>	<b>Percentage</b>
ISRNM (Gold standard)	83	56.8	57	54.8	26	61.9
MUAC	52	35.6	41	39.4	11	26.2
SGA	114	78.1	81	77.9	33	78.6
DMS	91	62.3	67	64.4	24	57.1
MIS	79	54.1	58	55.8	21	50
NPCR	97	66.4	69	66.3	28	66.7
Albumin	90	61.6	63	60.6	27	64.3
SGA + NPCR	134	91.8	94	90.4	40	95.2
DMS + NPCR	119	81.5	85	81.7	34	81
MIS + NPCR	111	76	79	76	32	76.2

DMS – Dialysis Malnutrition Score , ISRNM- International Society of Renal Nutrition and Metabolism, MIS – Malnutrition Inflammation Score, MUAC- Mid upper arm circumference, N PCR – Normalised Protein Catabolic Rate, SGA – Subjective Global Assessment