

Supplementary Table 2: Change in Acid-Base Parameters with Haemodialysis

	pH	N = 107 (%)	Mean
Pre-HD	<7.35	60 (56.07)	7.33±0.07
	7.35 – 7.45	44 (41)	
	>7.45	03 (2.8)	
	pH	N = 106 (%)	Mean
Post-HD	<7.35	18 (16.9)	7.40±0.068
	7.35 – 7.45	65 (61.3)	
	>7.45	23 (21.7)	
	Bicarbonate (mEq/l)	N = 107 (%)	Mean
Pre-HD	<18	42 (39.25)	18.60±2.86
	18-22	52 (48.59)	
	>22	13 (12.15)	
Post-HD	<22	09 (8.4)	26.14±2.94
	22-24	15 (14.01)	
	>24	83 (77.57)	
	pCO2 mEq/l	N = 107 (%)	Mean (mmHg)
Pre-HD	<35	61 (57)	35.69±6.80
	35 -45	41 (38.31)	
	>45	5 (4.6)	
Post-HD	<30	18 (16.82)	45.25±9.57
	30 -45	51 (47.66)	
	>45	38 (35.51)	

Supplementary Table 3: Protein Intake, Serum Albumin & Lean Body Mass

Protein Intake (Diet Diary) gm/kg/day	N = 107 (%)	Mean
<0.6	99 (92.52)	0.45±0.11 g/kg/day
0.6 – 0.8	7 (6.54%)	
0.8 – 1	1 (0.9)	
>1	0 (0)	
Protein Intake (nPCR) gm/kg/day	N = 107 (%)	
<0.6	36 (33.6)	0.73±0.23 g/kg/day
0.6 – 0.8	37 (34.6)	
0.8 – 1	19 (17.8)	
>1	15 (14)	
Serum Albumin (gm/dl)	N = 107 (%)	Mean (gm/dl)
<3.5	18 (16.82)	
3.5 – 5	85 (79.43)	
>5	4 (4.67)	
Lean Body mass (kg)	N = 107 (%)	Mean
<36	86 (80.37)	31.96±5.49
36-42	15 (14)	
>42	6 (5.6)	