

Supplementary Table 2. Dosing schedule for ravalizumab or crovalimab

Ravalizumab			
Body weight	Induction		Maintenance (2 weeks after induction)
	Intravenous (IV) infusion¹		IV infusion¹
≥100 kg	3000 mg once		3600 mg q 8 wk
≥60 kg to <100 kg	2700 mg once		3300 mg q 8 wk
≥40 kg to <60 kg	2400 mg once		3000 mg q 8 wk
≥30 to <40 kg	1200 mg once		2100 mg q 8 wk
≥20 to <30 kg	900 mg once		2100 mg q 8 wk
≥10 to <20 kg	600 mg once		600 mg q 4 wk
≥5 to <10 kg	600 mg once		300 mg q 4 wk
Crovalimab²			
Body weight	Induction		Maintenance
	Day 1: IV infusion	Days 2, 8, 15, 22: subcutaneously	Day 29, then q 4 wk: subcutaneously
≥100 kg	1500 mg	340 mg	1020 mg
≥40 kg to <100 kg	1000 mg	340 mg	680 mg

¹The drug is diluted to a concentration of 50 mg/mL for 3 mL and 11 mL vials and 5 mg/mL for 30 mL vials

²Currently licensed (elsewhere) for use only above 12 years of age and 40 kg body weight